



SAMPLE TYPE OR NO. *	DEPTH																			
	490-510ft C	789-809ft C	1090-1110ft C	1214-1234ft C	1506-1526ft C	1778-1798ft C	1931-1951ft C	3158-3168ft C	4011-4021ft C	5161-5171ft C	6327-6247ft C	7544-7550ft C	8690-8707ft C	9156-9176ft C	9641-9656ft C	10045-10065ft C				
FOSSIL NAMES																				
Proteacidites biporus																				
P. callosus																				
P. crassus	.																			
P. differentipollis	.	.		.																
P. dierama ms							.													
P. dilwynensis-grandis complex	.	.	.	.																
P. incurvatus		.	.	.																
P. kopiensis		.																		
P. latrobensis	.	.																		
P. leightonii	.			C																
P. nasus		.																		
P. obscurus	.	.	.	.																
P. ornatus	.																			
P. otwayensis ms							.													
P. pachypolus	.	.																		
P. pseudomoides																				
P. recavus	.																			
P. rectomarginis																				
P. rectus		.	.																	
P. reticulatus	.			C																
P. reticuloconcavus ms							.													
P. reticulosabratus		.		C																
P. retiformis				.	.															
P. rugulatus	.																			
P. scitus																				
P. tenuixinus																				
P. tuberculatus																				
P. tuberculiformis																				
P. tuberculotumulatus ms		.																		
P. spp. undescribed	.	.	.	.	.	.						C								
Protophloxypinus spp. R																				
Pseudowinterapollis calathus																				
P. cranwellae							.													
P. wahoensis							.													
Pyrolobospora reticulata																				
Reticulatisporites pudens									.											
Retistephanocolpites nixonii ms		.	.																	
Riccia boxatus ms																				
Rogalskaisporites cf canalis										.										
Rotverrusporites stellatus ms				.	.															
Rouseisporites reticulatus						.	.	.	.											
R. simplex																				
Rugulatisporites mallatus complex	.		.	.	.															
Santalumidites cainozoicus	.			C																
Sapotaceoidapollenites rotundus																				
Schizaea digitatoides					.															
Schizocolpus marlinensis																				
S. rarus ms																				
Selagosporis sp.																				
Spinizonocolpites prominatus																				
Stereisporites antiquisporites			.	.	.	.	.	.	.	.	.	.	.	.	.					
S. australis f. crassa			.			.	.													
S. cf pocockii							.	.	.	.	.	.	.	.	.					
S. punctatus ms			.	.	.	.														
S. regium ms						.	.													
S. spp. indeterminate	.	.	.	.	.	.														

\* C=CORE S=SIDEWALL CORE  
T=CUTTINGS J=JUNK BASKET

R = REWORKED SP.  
C = CONTAMINANT

DEPT. PRIMARY INDUSTRIES  
 PETROLEUM DEVELOPMENT  
 THIS DOCUMENT HAS BEEN  
 SCANNED  
 ON OR BEFORE MARCH 2000