

Palynology Report by L.E. Stover + A.D. Partridge with revisions



ASIN GIPPSLAND

DATE _____

ELL NAME ALBACORE - 1

ELEVATION + 99 feet

AGE	PALYNOLOGIC ZONES	HIGHEST DATA				LOWEST DATA					
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
Eocene	<u>P. tuberculatus</u>										
	<u>U. N. asperus</u>										
	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>										
	<u>P. asperopolus</u>										
	<u>U. M. diversus</u>										
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>										
Paleocene	<u>U. L. balmei</u>										
	<u>L. L. balmei</u>	9164	0				9196	0			
	<u>T. longus</u>	9422	2	9610	1		9956	1			
Cretaceous	<u>T. lilliei</u>	10,020	1				10,574	1			
	<u>N. senectus</u>	9120									
	<u>C. trip./T. pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											

COMMENTS: Trithyrodinium evittii Dinoflagellate Zone: 9164(0) to 9196(0)

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: L.E.S./A.D.P.

DATE June 1971, Dec. 1971

DATA REVISED BY: A.D.P.

DATE Jan. 1975.