

| DEPTH (metres) | CONTINENTAL SLOPE | SHELF EDGE → MID SHELF | MID-INNER SHELF |
|----------------|----------------------------------|------------------------|-----------------|
| | SIDEWALL CORE Depth in metres | | |
| 519.0 | <i>Oridorsalis tenera</i> | | |
| 538.0 | <i>Reophax</i> spp. | | |
| 555.0 | <i>Rhabdammina abyssorum</i> | | |
| 570.0 | <i>Discammina compressa</i> | | |
| 590.0 | <i>Martinottiella communis</i> | | |
| 605.0 | <i>Amnobaeculites calcaratus</i> | | |
| 627.5 | <i>Bathysiphon</i> spp. | | |
| 640.0 | <i>Stilostomella antillea</i> | | |
| 657.0 | <i>Bulimina marginata</i> | | |
| 674.0 | <i>Ammodiscus incertus</i> | | |
| 691.0 | <i>Brachisiphon corbiformis</i> | | |
| 708.0 | <i>Karreriella bradyi</i> | | |
| 727.0 | <i>Siphovigerina proboscidea</i> | | |
| 748.0 | <i>Gyroidinoides zealandica</i> | | |
| 758.0 | <i>Nonionella</i> spp. | | |
| 776.0 | <i>Pleurostomella tenera</i> | | |
| 790.0 | <i>"Pycnammina" spp.</i> | | |
| 810.0 | <i>Amnobaeculites</i> sp. | | |
| 827.0 | <i>Purgø Terebra</i> | | |
| 844.0 | | | |
| 861.0 | | | |
| 878.0 | | | |
| 892.5 | | | |
| 912.0 | | | |
| 929.0 | | | |
| 945.0 | | | |
| 963.0 | | | |
| 980.0 | | | |
| 993.0 | | | |
| 1013.0 | | | |
| 1150.0 | indet | | |
| 1200.0 | | | |
| 1250.0 | | | |
| 1300.0 | | | |
| 1350.0 | | | |
| 1400.0 | indet | | |
| 1450.0 | | | |
| 1500.0 | | | |
| 1550.0 | | | |
| 1600.0 | | | |
| 1650.0 | | | |
| 1700.0 | | | |
| 1750.0 | indet | | |
| 1800.0 | | | |
| 1850.0 | | | |
| 1900.0 | | | |
| 1950.0 | | | |
| 2000.0 | | | |
| 2025.0 | | | |
| 2050.0 | | | |
| 2070.0 | | | |
| 2090.0 | | | |
| 2115.0 | | | |

KEY: * = <20 specimens
 x = >20 specimens
 D = Dominant >60% specimens
 { = environmentally misplaced specimens
 ~~~~~ = definite hiatus  
 indet = specifically indeterminate due to diagenesis.

TABLE 3: DISTRIBUTION OF SELECTED BENTHONIC FORAMINIFERA IN BASKET # 1.

David Taylor, 9/8/1983.