



PE990316

BASIN GIPPSLAND

DATE _____

WELL NAME BATFISH -1ELEVATION +31 FEET

AGE	PALYNOLOGIC ZONES	HIGHEST DATA					LOWEST DATA				
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
IG-IO. Eocene	<u>P. tuberculatus</u>	4765	1				4768	1			
	<u>U. N. asperus</u>										
	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>										
	<u>P. asperopolus</u>	5004	1				5698	1			
	<u>U. M. diversus</u>	5956	1				6102	1			
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>										
PALEOCENE	<u>U. L. balmei</u>										
	<u>L. L. balmei</u>	6309	1				6740	1			
	<u>T. longus</u>	7332	1				8040	1			
CRETACEOUS	<u>T. lilliei</u>	8402	1				9067	1			
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											
	<u>T.D.</u>	9761									

COMMENTS:

Eisenackia crassitabulata Dino-flagellate Zone 6309' (2)None of the Wetzeliella Zones can be indentified within the Flounder Fm. in this well.

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
- 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
- 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
- 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
- 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: LES/ADPDATE June 1971 ; Dec 1971.DATA REVISED BY: ADPDATE Jan 1975.