



PE990406

BASIN GIPPSLAND

BY David J. Taylor

WELL NAME FLOUNDER-4

DATE 5/2/73

ELEV. +32

Foram Zonules

		Highest Data	Quality	2 Way Time	Lowest Data	Quality	2 Way Time
MIOCENE	A	Alternate					
	B	Alternate					
	C	Alternate					
	D	3258	1		5160	1	
	D 1	Alternate					
	D	5500	2		5774	1	
	D 2	Alternate	5774	1			
	E	6150	1		6200	1	
	E	Alternate					
	F	6230	0		6335	1	
	F	Alternate					
OLIGOCENE	G	Alternate					
	H 1	Alternate					
	H 2	Alternate					
	I 1	Alternate					
	I 2	Alternate					
	J	6345	1		6345	1	
ECC.	J 1	Alternate					
	J 2	Alternate					
K	Alternate						
Pre K *	6610	1		6748	1		

COMMENTS: * Equals Lower Eocene determined on planktonic fauna, on benthonic foraminifera the Lower Eocene interval is probably between 6460 - 6806 feet.

Note: If highest or lowest data is a 3 or 4, then an alternate 0, 1, 2 highest or lowest data will be filled in if control is available.

If a sample cannot be interpreted to be one zonule, as apart from the other, no entry should be made.

- 0 SWC or Core - Complete assemblage (very high confidence).
- 1 SWC or Core - Almost complete assemblage (high confidence).
- 2 SWC or Core - Close to zonule change but able to interpret (low confidence).
- 3 Cuttings - Complete assemblage (low confidence).
- 4 Cuttings - Incomplete assemblage, next to uninterpretable or SWC with depth suspicion (very low confidence).