



AGE	PALYNOLOGIC ZONES	HIGHEST DATA					LOWEST DATA				
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
OLIG-MIO.	<u>P. tuberculatus</u>	2421	0				2441	0			
	<u>U. N. asperus</u>										
EOCENE	<u>M. N. asperus</u>										
	<u>L. N. asperus</u>										
	<u>P. asperopolus</u>										
	<u>U. M. diversus</u>										
	<u>M. M. diversus</u>										
	<u>L. M. diversus</u>	2444.9	0				2516.5	1			
	<u>U. L. balmei</u>	2553.5	0				2636	1			
PALEOCENE	<u>L. L. balmei</u>										
	<u>T. longus</u>										
LATE CRETACEOUS	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											

COMMENTS: Sidewall core #16 from 2546m was barren of specific index fossils and could not be designated as L. M. diversus or U. L. balmei with surety. All depths in metres

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.  
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.  
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.  
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.  
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: H.E. Stacy DATE January 22, 1979

DATA REVISED BY: \_\_\_\_\_ DATE \_\_\_\_\_