

Core Sample ID	Sample Depth (m)																																								Notes			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40				
Belfast No.4 c.AD 5581-65'																																											Insect	
c.AU 5065-73'																																												Paranoxa
c.AW 5344-54'		+				+														+																						Indet		
c.AX 5501-21'																																												
Heywood No.10 c.BP 4688-95'																							+				+																	
c.BR 4809-88'			R																																									
c.BS 4920 85'																																												
c.BW 5244-30'																																												
c.BX 4189-500'	Blacked out row																																								III			
Narrawaturk No.2 c.BE 4715-30'																																												
c.BF,BG 4973-80'																																												
c.BH 5147-50'																																												
c.BI 5327-36'																																												
Portland No.3 c.BP 5273-86'																																												
c.BQ 5467-76'																																												
Latrobe No.1 c.CY 1400-15'																																												
c.DB 1627-51'																																												
c.DC 1735-39'																																											III	
Ecklin No.3 c.AO 2142-44'																																												
c.AR 2472-90'		+																																								Spec obs		
c.AS 2581-68'																																										Insect		

Table 1. Distribution of selected spore, pollen, and microplankton species in core samples from Yangery No.1, Laang No.1, Belfast No.4, Heywood No.10, Narrawaturk No.2, Portland No.3, Latrobe No.1, and Ecklin No.3 bores.

+ - species present
 cf - specimens similar to, but not identical with, a particular species
 R - reworked spores